

# HOUSE OF WISDOM: JOURNAL ON LIBRARY AND INFORMATION SCIENCES

ISSN 3089-6002. Published by Universitas Muhammadiyah Sidoarjo  
Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC-BY).

## Adaptive Osmotic Filtration Mechanisms in Sap-Feeding Insects

Naghm Abdulghani mohammed <sup>1</sup>, Fadel Abbas Qader <sup>2\*</sup>

college of education for women, Kirkuk university, Iraq

college of medicinal and industrial plants, Kirkuk university, Iraq

Email : [naghm.abdoalghany@gmail.com](mailto:naghm.abdoalghany@gmail.com)

**Abstract. General Background:** Sap-feeding insects consume diets rich in water but poor in essential nutrients, creating major physiological challenges.

**Specific Background:**

Many Hemipteran insects possess a specialized digestive modification known as the filter chamber that facilitates sap processing.

**Knowledge Gap:** Although its anatomical presence is well documented, an integrated understanding of its functional mechanisms, molecular basis, and evolutionary diversification remains limited.

**Aims:** This review synthesizes current

anatomical, physiological, and molecular findings on the insect filter chamber, with emphasis on digestion, absorption, and evolutionary specialization.

**Results:**

Evidence shows that the filter chamber enables rapid osmotic transfer of excess water and sugars from the foregut-midgut directly to the hindgut, supported by aquaporins and sugar transporters, thereby optimizing nutrient retention and metabolic efficiency. Structural complexity varies across taxa, reflecting host-plant adaptation.

**Novelty:** The article integrates classical morphology with recent molecular insights to present the filter chamber as a dynamic regulatory system rather than a passive structure.

**Implications:** Understanding filter chamber function offers prospects for targeted pest management strategies through disruption of specific transport pathways while reducing reliance on broad-spectrum insecticides.

### Highlights:

- The filter chamber regulates water and sugar balance during sap feeding.
- Structural variation reflects evolutionary adaptation to host plants.
- Transport proteins present potential targets for selective pest control.

### Keywords:

Filter Chamber; Sap-Feeding Insects; Osmotic Regulation; Aquaporins; Digestive Adaptation

## Introduction

### Taxonomic Position of Insects in the Animal Kingdom

Insects belong to the class **Insecta**, which falls under the subphylum **Hexapoda** within the phylum **Arthropoda**, a major division of the kingdom **Animalia** [1, 39]. Insects form the most abundant and diverse animal group, with more than one million described species [2].

The class Insecta includes several orders, such as:

- **Coleoptera** (beetles and weevils) [36, 38]
- **Lepidoptera** (butterflies and moths)
- **Hymenoptera** (ants, bees, and wasps)

**Commented [j1]:** Pendahuluan artikel telah memuat latar belakang umum yang jelas mengenai keragaman serangga dan tantangan nutrisi pada serangga pemakan cairan. State of the art ditunjukkan melalui pemaparan penelitian terdahulu tentang sistem pencernaan dan filter chamber. Kesenjangan penelitian juga mulai terlihat pada kebutuhan integrasi antara pendekatan morfologi dan molekuler. Tujuan penelitian disampaikan dengan cukup tegas, yakni meninjau fungsi dan modifikasi evolusioner filter chamber. Unsur kebaruan muncul dari upaya menggabungkan temuan klasik dan studi modern, meskipun dapat ditegaskan kembali secara lebih eksplisit.

# HOUSE OF WISDOM: JOURNAL ON LIBRARY AND INFORMATION SCIENCES

ISSN 3089-6002. Published by Universitas Muhammadiyah Sidoarjo  
Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC-BY).

- **Hemiptera** (bugs, aphids, whiteflies)
- **Orthoptera** (grasshoppers, locusts, crickets)
- **Diptera** (mosquitoes and flies) [3, 4]

## 2. Feeding Types and Their Relationship to Mouthpart Structure in Insects

Insect feeding habits vary widely, and insect mouthparts have adapted to feed on a wide variety of food sources [5]. Some of the most notable changes to insect mouthparts:

- **Biting mouthparts**: such as in beetles and grasshoppers.
- **very long tube-like mouthparts** which penetrate plant or animal tissue as in aphids and mosquitoes.
- **Siphoning mouthparts**: adapted to nectar-feeding, like in butterflies.
- **Sponging-lapping**: for soaking up freely available liquid food, like houseflies.
- **Chewing—sucking mouthparts**: suited to gathering and absorbing nectar, such as those in bees [6].

## Method

### 3. Feeding Patterns in Insects

#### 3.1 Herbivorous Insects

Most insect species are herbivorous insects [7]. The various parts of plants including fruits, leaves, stems, roots, nectar and sap are their food [8]. These insects encounter various obstacles, such as plant mechanical and chemical protection and a poor protein level of sap of plants [37]. As a result, they developed an elaborated category of specialized physiological structures (filter chamber [9], variation in gut length according to dietary quality [35].

#### 3.2 Carnivorous Insects

This set includes carnivorous insects such as lacewings and parasitic species, including some wasps [10]. They have protein and lipid-rich diet, strong chewing or piercing sucking mouth parts [11].

#### 3.3 Omnivorous Insects

These insects show a considerable dietary flexibility and ability to use either plant or animal food sources, like toxic plants for cockroaches or ants [12]. Their alimentary canal is generally less modified than in strictly herbivorous or carnivorous species [13].

**Commented [j2]:** Metode yang digunakan dalam artikel ini sudah sesuai dengan karakteristik penelitian berupa artikel tinjauan. Sumber data berasal dari literatur ilmiah yang relevan dan kredibel, mencakup buku referensi serta jurnal internasional bereputasi. Teknik pengumpulan data melalui penelusuran pustaka telah sejalan dengan tujuan kajian konseptual. Instrumen utama berupa analisis dokumen juga tepat digunakan. Teknik analisis data dilakukan secara naratif dan sintesis tematik, yang sesuai dengan pendekatan review, meskipun penjelasan tentang kriteria pemilihan literatur dapat diperjelas.

## HOUSE OF WISDOM: JOURNAL ON LIBRARY AND INFORMATION SCIENCES

ISSN 3089-6002. Published by Universitas Muhammadiyah Sidoarjo  
Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC-BY).

### 3.4 Saprophagous Insects

They feed on decomposing organic matter - carrion beetles and certain flies' larvae. They are significant decomposers of organic matters and nutrient recycler [14]. Their digestion frequently relies on the gut microorganisms and microbes participate in degradation of complex molecules such as keratin and cellulose [15, 16].

## 4. Herbivorous Insect Digestive Structures

Insects have a three-region digestive system:

### 4.1 Foregut

Consists of the pharynx, esophagus, crop and proventriculus. It serves for collecting the food, for its temporary retention and also has a role in initial mechanical processing [17].

### 4.2 Midgut

This is the major area for digestion and absorption. It produces digestive enzymes which aid in the digestion and absorption of nutrients [18].

### 4.3 Hindgut

Functions of the kidney include absorption of fluids and salts as well as excretion of metabolites [19].

## 5. Excretory Products of Sap-Feeding Insects

Homopterans, such as aphids generate honeydew, a large amount of sugary liquid waste [20]. Honeydew is mainly composed of non-metabolized sugars, glucose, sucrose and fructose including water and represents a significant food resource for various insects like ants [21].

## Result

### 6. The Filter Chamber in Insects

The filter chamber is a separate anatomical structure (Figure 1) [31], where the crop (foregut-hindgut juncture) wraps closely with thin membrane around the midgut, encloses an open cavity surrounded by basal membrane folds [22, 23].

This structure is found mainly in insects of the order **Hemiptera** [40] and in some members of the order **Thysanoptera** [24].

**Commented [j3]:** Pembahasan dalam artikel telah menguraikan hasil kajian secara relatif rinci dan berjenjang, mulai dari aspek morfologi hingga mekanisme molekul filter chamber. Penulis secara konsisten mengaitkan temuan dengan teori dan hasil penelitian sebelumnya, sehingga argumen yang dibangun terasa logis dan kontekstual. Integrasi antara data anatomi klasik dan studi modern tentang transporter serta aquaporin memperkaya kedalaman pembahasan. Keterkaitan antara struktur, fungsi, dan adaptasi evolusioner juga dijelaskan dengan cukup baik. Namun, pada beberapa bagian, pembahasan masih cenderung deskriptif dan dapat diperdalam dengan analisis komparatif yang lebih kritis antar kelompok serangga.

## HOUSE OF WISDOM: JOURNAL ON LIBRARY AND INFORMATION SCIENCES

ISSN 3089-6002. Published by Universitas Muhammadiyah Sidoarjo  
Copyright © Author(s). This is an open-access article distributed under the  
terms of the Creative Commons Attribution License (CC-BY).




Fig. 10. A. Alimentary canal of *Triecphora vulnerata* (after LICENT, 1912). B. Transverse section of filtercomplex showing muscular sheath (ms), filterchamber (fic), stomach (st), intestine (i), and MALPIGHIAN tubules (Mt). C. Simplified diagram of the filtercomplex available for all cicadoids summarized in Table 5. In some cicadooids the posterior part of the midgut forms a simple loop within the filterchamber whereas in other ones it is coiled to various degrees.

Figure 1 [31]

### 7. Functional and Evolutionary Modifications of the Filter Chamber and Their Relation to Digestion and Absorption

This osmotic filtration is essential for concentrating sap ingested [25]. The partial concentration of plant sap leads to fluorescent xylem and the xylem encourages counterflow due to proximity with the foregut. Surplus water and monosaccharides (glucose and fructose) are transported osmotically from the lumen of the foregut-midgut directly into the hindgut [26].

Such transport permits a slow transit of nutrients, like amino acids, through the midgut thereby promoting efficient digestion and absorption [27]. Furthermore, the very quick removal of excess sugars and water over the rectum avoids dilution loss of digested enzyme and saves energy [28].

Studies have found that membranes of filter chamber cells contain highly developed aquaporin water channels and sugar transporters, to control the movement of solutes [29,

## HOUSE OF WISDOM: JOURNAL ON LIBRARY AND INFORMATION SCIENCES

ISSN 3089-6002. Published by Universitas Muhammadiyah Sidoarjo  
Copyright © Author(s). This is an open-access article distributed under the  
terms of the Creative Commons Attribution License (CC-BY).

30].

The nature of the filter chamber is different among insects. It is rather primitive in aphids (Aphidoidea) [31], but more advanced and morphologically complex in whiteflies (Aleyrodidae) and scale insects (Coccoidea), which allows them to feed on certain categories of plant sap [32, 33].

### Conclusions

1. The filter chamber represents a remarkable evolutionary adaptation that enables insects to overcome the challenges of feeding on a nutrient-poor, water-rich diet.
2. Its function relies primarily on a precise physio-chemical osmotic filtration mechanism supported by a specialized anatomical configuration.
3. Recent molecular studies have revealed significant complexity in the processes governing water and solute transport within this organ.
4. Variation in filter chamber structure across hemipteran insects reflects evolutionary specialization for adapting to different host plants.

**Commented [j4]:** Kesimpulan pada artikel ini pada dasarnya sudah menjawab tujuan penelitian yang ditetapkan sejak awal. Penulis berhasil merangkum temuan utama mengenai peran dan karakteristik filter chamber sebagai adaptasi fisiologis serangga pemakan cairan. Hubungan antara tujuan kajian, yaitu memahami fungsi, mekanisme, dan variasi evolusioner filter chamber, dengan simpulan yang disajikan terlihat cukup jelas. Kesimpulan juga menegaskan kontribusi artikel sebagai tinjauan komprehensif, bukan sekadar deskripsi, sehingga pembaca memperoleh gambaran menyeluruh tentang signifikansi topik yang dibahas. Meski demikian, simpulan akan terasa lebih kuat apabila disertai penegasan kembali implikasi ilmiah dan aplikatif secara lebih eksplisit.

### Recommendations

1. Transcriptomic and proteomic studies of the cells lining the filter chamber to clarify gene and protein networks that control transport activities are urgently required.
2. New live imaging recording techniques, such as confocal microscopy should be used in this real time transport study of water and solute.
3. Exploration of the possibilities to attack certain transporter proteins in the filter chamber (sugar specific transporters or aquaporins, for example) may contribute to finding new selective means of control for sap-consuming pests, decreasing dependence from broad-spectrum insecticides [34].
4. Gut microbiota functions in digestion/nutrient utilization of insects with a filter chamber, and this is especially true for sap feeding as mandated by studies.

### Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of and other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

### Conflict of Interest Statement

## HOUSE OF WISDOM: JOURNAL ON LIBRARY AND INFORMATION SCIENCES

ISSN 3089-6002. Published by Universitas Muhammadiyah Sidoarjo  
Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC-BY).

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

### References

- [1] Z.-Q. Zhang, "Phylum Arthropoda," in *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness (Addenda 2013)*, Z.-Q. Zhang, Ed. Auckland, New Zealand: Magnolia Press, 2013, pp. 1–82.
- [2] N. E. Stork, "How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth?" *Annual Review of Entomology*, vol. 63, pp. 31–45, 2018.
- [3] D. Grimaldi and M. S. Engel, *Evolution of the Insects*. Cambridge, UK: Cambridge University Press, 2005.
- [4] P. J. Gullan and P. S. Cranston, *The Insects: An Outline of Entomology*, 5th ed. Oxford, UK: Wiley-Blackwell, 2014.
- [5] R. F. Chapman, *The Insects: Structure and Function*, 5th ed. Cambridge, UK: Cambridge University Press, 2013.
- [6] F. Karolyi, J. F. Colville, S. Handschuh, B. D. Metscher, and H. W. Krenn, "One Proboscis, Two Tasks: Adaptations to Blood-Feeding and Nectar-Extracting in Long-Proboscid Horse Flies," *Arthropod Structure & Development*, vol. 41, no. 3, pp. 275–281, 2012.
- [7] L. M. Schoonhoven, J. J. A. van Loon, and M. Dicke, *Insect-Plant Biology*, 2nd ed. Oxford, UK: Oxford University Press, 2005.
- [8] A. R. War, G. K. Taggar, M. Y. War, and B. Hussain, "Impact of Climate Change on Insect Pests, Plant Chemical Ecology, and Tritrophic Interactions," in *Sustainability in Plant and Crop Protection*. Cham, Switzerland: Springer, 2016.
- [9] A. E. Douglas, "Phloem-Sap Feeding by Animals: Problems and Solutions," *Journal of Experimental Botany*, vol. 57, no. 4, pp. 747–754, 2006.
- [10] M. A. Jervis, *Insects as Natural Enemies: A Practical Perspective*. Dordrecht, The Netherlands: Springer, 2005.
- [11] L. I. Gilbert and S. S. Gill, Eds., *Insect Control: Biological and Synthetic Agents*. London, UK: Academic Press, 2010.
- [12] M. Coll and M. Guershon, "Omnivory in Terrestrial Arthropods: Mixing Plant and Prey Diets," *Annual Review of Entomology*, vol. 47, pp. 267–297, 2002.
- [13] W. R. Terra and C. Ferreira, "Basic Biology of Insect Midgut and Its Associated Microbiota," in *Insect Gut Microbiome and Its Applications*. Cham, Switzerland: Springer, 2020.
- [14] M. E. Benbow, J. K. Tomberlin, and A. M. Tarone, Eds., *Carrion Ecology, Evolution, and Their Applications*. Boca Raton, FL, USA: CRC Press, 2015.
- [15] P. Engel and N. A. Moran, "The Gut Microbiota of Insects: Diversity in Structure and Function," *FEMS Microbiology Reviews*, vol. 37, no. 5, pp. 699–735, 2013.
- [16] A. E. Douglas, "Multi-Organismal Insects: Diversity and Function of Resident Microorganisms," *Annual Review of Entomology*, vol. 60, pp. 17–34, 2015.
- [17] J. L. Nation, *Insect Physiology and Biochemistry*, 3rd ed. Boca Raton, FL, USA: CRC Press, 2016.
- [18] W. R. Terra and C. Ferreira, "Biochemistry and Molecular Biology of Digestion," in *Insect Molecular Biology and Biochemistry*. London, UK: Academic Press, 2012, pp. 365–418.
- [19] J. A. Dow, "The Essential Roles of Metal Ions in Insect Homeostasis and Physiology," *Current Opinion in Insect Science*, vol. 23, pp. 43–50, 2017.

## HOUSE OF WISDOM: JOURNAL ON LIBRARY AND INFORMATION SCIENCES

ISSN 3089-6002. Published by Universitas Muhammadiyah Sidoarjo  
Copyright © Author(s). This is an open-access article distributed under the  
terms of the Creative Commons Attribution License (CC-BY).

- [20] M. K. Fischer and A. W. Shingleton, "Host Plant and Ants Influence the Honeydew Sugar Composition of Aphids," *Functional Ecology*, vol. 15, no. 4, pp. 544–550, 2001.
- [21] B. Stadler and A. F. Dixon, "Ecology and Evolution of Aphid-Ant Interactions," *Annual Review of Ecology, Evolution, and Systematics*, vol. 36, pp. 345–372, 2005.
- [22] A. J. P. Goodchild, "Evolution of the Alimentary Canal in the Hemiptera," *Biological Reviews*, vol. 41, no. 1, pp. 97–140, 1966.
- [23] J. Pompon, D. Quiring, C. Goyer, P. Giordanengo, and Y. Pelletier, "A Phloem-Sap Feeder Mixes Phloem and Xylem Sap to Regulate Osmotic Potential," *Journal of Insect Physiology*, vol. 57, no. 9, pp. 1317–1322, 2011.
- [24] D. E. Ullman and D. L. McLean, "The Alimentary Canal of *Frankliniella occidentalis*," *International Journal of Insect Morphology and Embryology*, vol. 17, no. 2, pp. 125–134, 1988.
- [25] S. Caccia, M. Casartelli, and G. Tettamanti, "The Amazing Complexity of Insect Midgut Cells," *Cell and Tissue Research*, vol. 377, no. 3, pp. 505–525, 2019.
- [26] D. R. Price and J. A. Gatehouse, "RNAi-Mediated Crop Protection Against Insects," *Trends in Biotechnology*, vol. 26, no. 7, pp. 393–400, 2014.
- [27] D. A. Ashford, W. A. Smith, and A. E. Douglas, "Living on a High Sugar Diet," *Journal of Insect Physiology*, vol. 46, no. 3, pp. 335–341, 2000.
- [28] T. L. Wilkinson and A. E. Douglas, "Phloem Amino Acids and Host Plant Range," *Entomologia Experimentalis et Applicata*, vol. 106, no. 2, pp. 103–113, 2003.
- [29] G. D. Ewart and A. J. Howells, "ABC Transporters Involved in Transport of Eye Pigment Precursors," *Methods in Enzymology*, vol. 292, pp. 213–224, 1998.
- [30] S. Kikuta et al., "Aquaporin Water Channel AgAQP1," *Journal of Insect Physiology*, vol. 58, no. 6, pp. 829–835, 2012.
- [31] M. B. Ponsen, *The Digestive System of Subsaltusaphis ornata*. Wageningen, The Netherlands: Agricultural University Wageningen, 1979.
- [32] J. M. Cicero, J. K. Brown, and P. D. Roberts, "The Digestive System of the Whitefly *Bemisia tabaci*," *International Journal of Insect Morphology and Embryology*, vol. 24, no. 3, pp. 295–306, 1995.
- [33] I. Foldi, "The Scale Cover," in *Soft Scale Insects: Their Biology, Natural Enemies and Control*, vol. 7A. Amsterdam, The Netherlands: Elsevier, 1997.
- [34] I. K. Singh and S. Singh, "RNA Interference: A Novel Source of Resistance to Combat Whitefly," in *RNAi for Plant Improvement and Protection*. Wallingford, UK: CABI, 2021.
- [35] D. A. Ashford, W. A. Smith, and A. E. Douglas, "Living on a High Sugar Diet," *Journal of Insect Physiology*, vol. 46, no. 3, pp. 335–341, 2000.
- [36] A. M. Jamal, A. A. Haidar, and M. O. Mohammed, "Extraction and Evaluation of the Effectiveness of Chitosan as Insecticides," *International Journal of Agriculture and Animal Production*, vol. 4, no. 6, pp. 16–24, 2024.
- [37] T. S. A. Al-Hayali, O. T. H. A. Alhadithy, and F. A. Qader, "Toxic Efficiency of Fumigation Using Two Essential Oils," *Indian Journal of Agricultural Research*, vol. 59, no. 6, pp. 988–992, 2025.
- [38] S. H. A. K. Al-Hamawandy and S. M. Shahrabani, "Integrated Management of the Oriental Citrus Mite," *Kirkuk University Journal for Agricultural Sciences*, vol. 15, no. 1, pp. 47–53, 2024.
- [39] A. M. Mahmoud and S. S. Hamad, "Molecular Study of Head Lice Among Primary School Children," *Kirkuk Journal of Science*, vol. 20, pp. 43–47, 2025.
- [40] A. D. Yates and A. Michel, "Mechanisms of Aphid Adaptation to Host Plant Resistance," *Current Opinion in Insect Science*, vol. 26, pp. 41–49, 2018..